

NANO ENGINEERED

Growing with

FERTION



MCZ Metalin 524

**Microelement application for horticultural and agricultural use.
Liquid micronutrient for foliar and/or soil application.**

5 Litre

Gross weight: 8.05 kg Nett weight: 7.75 kg
SG 1.55 at 20 °C

Highly concentrated | Bio available

REGISTERED PLANT NUTRIENT CONTENT

Copper as Cu	110 g/l (73.8 g/kg)
Manganese as Mn	330 g/l (221.5 g/kg)
Zinc as Zn	84 g/l (56.4 g/kg)

Registered and Manufactured by:

Kimleigh Chemicals SA (Pty) Ltd
Reg. No. 1974/00531/07

Tel: +27 18 293 1028 Fax: +27 18 294 4079
11 Jasper van der Westhuizen Street, Potchefstroom, 2531

www.kimleigh.com



KIMLEIGH
CHEMICALS SA (PTY) LTD
INNOVATIVE CHEMISTRY



Fertilizer Group 2
Reg No B5184 Act 36 of 1947

Recommendations:

CROP	DOSAGE/APPLICATION*	WATER RATE	TIMING*
Wheat Maize Peas Grapes Onions Potatoes Leaf Vegetables Brassicas	1 ℓ / ha	200 ℓ / ha by tractor driven sprayer 100 000 ℓ / ha by irrigation system.	Repeat 2 – 3 times at 10 – 14 day intervals during crop lifecycle.

DIRECTIONS FOR USE

Mix the product thoroughly before use. Fill working solution tank with 75% water, agitate and pour in required product quantity. Top up working solution tank to capacity. Start spraying while agitating solution. Foliar/Soil: use enough water for full cover spray or to spray soil surface around the root zone. Care should be taken when mixing this product with other products in the same working solution tank and is entirely at the risk of the end-user.

PRECAUTIONS

Use only on crops as listed. Do not apply under extreme conditions such as rain, frost, very low and very high humidity. Wear PPE such as gloves and face shield when handling the product. Always clean application equipment before and after use.

WARNINGS

Harmful if swallowed. Do not breathe spray. Inhalation may cause irritation or damage to respiratory tract.

STORAGE

Keep in a cool, dry chemical store and avoid temperatures in excess of 40°C. Always seal used containers. Keep out of reach of children, livestock and uninformed persons.

*** Best practice: Product dosage should be determined by soil, leaf and/or sap analyses.**

NOTE: Adulteration mixing or dilution of this product and/or where it is used for something other than its intended purpose, will result in any warranty given by the Manufacturer declared as null and void.

Please contact your Agriculturist for specific crop application.

Shake well before use.

Skud deeglik voor gebruik.

